

# **Development of a Novel Coupled Simulation Tool to Study Ocean-Estuary Exchange**

Oliver B. Fringer  
473 Via Ortega, Room 187  
Dept. of Civil and Environmental Engineering  
Stanford University, Stanford, CA 94305  
phone: (650) 725-6878 fax: (650) 725-9720 email: [fringer@stanford.edu](mailto:fringer@stanford.edu)

Grant Number: N00014-08-1-0904  
<http://suntans.stanford.edu>

## **LONG-TERM GOALS**

The long-term goal of this project is to develop a coupled ocean-estuary simulation tool that will consist of the Regional Oceanic Modeling System (ROMS) and the nonhydrostatic, unstructured-grid SUNTANS model, in order to study the interaction of coastal regions with estuaries. The scientific goal will be to understand the dynamics of ocean-estuary exchange between San Francisco Bay and the coastal Pacific Ocean, with ROMS simulating the coastal Pacific Ocean, and SUNTANS simulating San Francisco Bay.

## **OBJECTIVES**

The primary objective of this project is to develop and implement a coupled hydrodynamics and sediment-transport simulation tool to study the complex three-dimensional, coupled ocean-estuary processes that influence exchange through the Golden Gate channel in San Francisco Bay. In order to understand the influence of regional processes in the coastal ocean and how they are coupled to local processes in San Francisco Bay, the project will employ a one-way coupled simulation tool, in which a model for San Francisco Bay is coupled to a regional model for the coastal Pacific Ocean. The SUNTANS model (Fringer et al., 2006a) will be employed to simulate flow in San Francisco Bay, and it will obtain boundary conditions at the coastal ocean from simulations of the California coastal current using ROMS (Shchepetkin and McWilliams, 2005). SUNTANS is well-suited to simulate the exchange flow at the Golden Gate because it employs an unstructured grid, which enables simulation domains that include the entire Bay while also resolving finescale flow features near the Golden Gate. Furthermore, because SUNTANS is a nonhydrostatic code, it will capture nonhydrostatic features of the exchange flow which may be important at high resolution. ROMS, on the other hand, was originally written to simulate regional flows like the California Coastal Current, and employs highly accurate methods to simulate the complex dynamics of upwelling fronts, jets, and squirts over seasonal time scales (Marchesiello, et al., 2003). Using a stand-alone SUNTANS, simulations will be performed in idealized domains to understand the fundamental mechanisms governing three-dimensional exchange. In the real, field-scale domain, one-way coupling will be used to understand how tidal and seasonal variability in the coastal currents as well as in the Bay influence the dynamics of ocean-estuary exchange through the Golden Gate channel.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 SEP 2008</b>		2. REPORT TYPE <b>Annual</b>		3. DATES COVERED <b>00-00-2008 to 00-00-2008</b>	
4. TITLE AND SUBTITLE <b>Development Of A Novel Coupled Simulation Tool To Study Ocean-Estuary Exchange</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Stanford University,Dept. of Civil and Environmental Engineering,473 Via Ortega, Room 187,Stanford,CA,94305</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>code 1 only</b>					
14. ABSTRACT <b>The long-term goal of this project is to develop a coupled ocean-estuary simulation tool that will consist of the Regional Oceanic Modeling System (ROMS) and the nonhydrostatic, unstructured-grid SUNTANS model, in order to study the interaction of coastal regions with estuaries. The scientific goal will be to understand the dynamics of ocean-estuary exchange between San Francisco Bay and the coastal Pacific Ocean, with ROMS simulating the coastal Pacific Ocean, and SUNTANS simulating San Francisco Bay.</b>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>4</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

## APPROACH

Simulations of San Francisco Bay will employ SUNTANS (Stanford Unstructured Nonhydrostatic Terrain-following Adaptive Navier-Stokes Simulator), a free-surface, nonhydrostatic, unstructured-grid, parallel coastal ocean and estuary simulation tool that solves the Navier-Stokes equations under the Boussinesq approximation (Fringer et al., 2006a). The formulation is based on the method outlined by Casulli & Walters (2000), in which the free-surface and vertical diffusion are discretized with the theta-method, which eliminates the Courant condition associated with fast free-surface waves and the elevated friction term associated with small vertical grid spacing at the free-surface and bottom boundary. For flows with extensive wetting and drying such as San Francisco Bay, advection of momentum is accomplished with the semi-Lagrangian advection scheme (Casulli and Cheng, 1992; Staniforth and Temperton, 1986), which ensures stability in the presence of cells that fill and empty with the tides.

To simulate the influence of coastal currents on exchange between San Francisco Bay and the coastal Pacific, the SUNTANS simulations will be nested within the ROMS model (Shchepetkin and McWilliams, 2005), which also has a nonhydrostatic module (Kanarska, et al., 2007). ROMS is discretized in horizontal curvilinear coordinates and has a generalized terrain-following vertical coordinate that is configured to enhance resolution near the sea surface, and the time-stepping algorithm is split-explicit.

The simulations will employ a one-way coupled ROMS-SUNTANS tool that has been developed and applied to simulate nonlinear internal tides on the Australian North West Shelf (Fringer et al., 2006b), and development of the associated two-way coupled tool is underway in collaboration with Prof. J. McWilliams at UCLA in order to study tide-current interactions in Monterey Bay as well as the South China Sea. The SUNTANS simulations will be one-way nested within simulations of the California Coastal Current (Marchesiello et al., 2003). These simulations will provide sea-surface height, velocity fields, and sediment, salinity and temperature at the offshore boundary for SUNTANS. Because the ROMS simulations of the coastal current are nontidal, tidal currents from the OTIS tidal model (Egbert et al., 1994) are included by adding them to the mesoscale currents provided by ROMS at the SUNTANS boundaries. In one-way nesting, the boundary conditions supplied by ROMS influence SUNTANS, yet it is assumed that the flowfield as computed by SUNTANS does not influence the ROMS simulation (see, e.g. Fringer et al., 2006c). The boundaries of the SUNTANS simulations will be placed in order to minimize their potential influence on the exchange plume as it evolves on the shelf offshore of the Golden Gate. As long as the plume that exits the SUNTANS boundary does not have the potential to reenter the Bay, then the boundary is far enough from the mouth of the Bay and one-way nesting is justified.

Because both SUNTANS and ROMS are well-established models and the one-way coupling framework has already been developed (Fringer et al., 2006b), a bulk of this project will focus on implementation of the models to understand detailed physics of ocean-estuary exchange. Yet while SUNTANS has successfully been applied to study an estuary with significant wetting and drying (Wang et al., 2008), advection of momentum in the presence of wetting and drying presents significant challenges when high grid resolution is employed. While accuracy of semi-Lagrangian advection schemes depends on the time-accuracy of the traceback as well as the spatial accuracy of the interpolation (Staniforth and Cote, 1991), typical implementations of semi-Lagrangian advection for estuarine simulations with coarse resolution employ first-order accurate spatial interpolation (Casulli

and Cattani, 1994) and ignore vertical advection of momentum (Zhao, 2007). While this does not incur problems when coarse resolution is employed, high-resolution requires implementation of vertical advection and higher-order interpolation in the presence of complex bathymetry. This is not a trivial task, since it is difficult to ensure stability in the implementation of higher-order interpolation schemes on unstructured, finite-volume grids, particularly with wetting and drying (Zhao, 2007). However, if the stencil is suitably chosen, stability can be ensured, as pointed out by Leonard (2002). In the proposed work we will explore stencils for the Lagrangian traceback that improve stability when second-order accurate interpolation is used.

## **WORK COMPLETED**

Funding began mid FY08 so no work has been completed to date.

## **IMPACT/APPLICATIONS**

The coupled simulation tool that will be developed for this project has the potential to form the basis for a wide range of modeling studies of other important coupled ocean-estuary systems.

## **RELATED PROJECTS**

The California Coastal Conservancy has recently funded the development of an open-source hydrodynamics and sediment transport model using SUNTANS to assess the impacts of marsh restoration projects on salt and sediment dynamics in San Francisco Bay. Fringer is involved in this project in collaboration with Profs. Mark Stacey and Zack Powell at U. C. Berkeley and Profs. Jeffrey Koseff and Stephen Monismith at Stanford University.

## **REFERENCES**

Casulli, V., and E. Cattani (1994), Stability, accuracy and efficiency of a semi-implicit method for three-dimensional shallow water flow, *Comp. Math. with Applic.*, 27, 99-112.

Casulli, V. and R.A. Walters (2000), An unstructured, three-dimensional model based on the shallow water equations, *Int. J. Numer. Meth. Fluids.*, 32, 331-348.

Egbert, G. D., A. F. Bennett, and M. G. G. Foreman (1994), TOPEX/ POSEIDON tides estimated using a global inverse model, *J. Geophys. Res.*, 99(C12), 24,821– 24,852.

Fringer, O. B., Gerritsen, M., and R. L. Street (2006a), An unstructured-grid, finite-volume, nonhydrostatic, parallel coastal ocean simulator, *Ocean Modelling*, 14 (3-4), 139-278.

Fringer, O. B., Gross, E. S., Meulenens, M., and G. N. Ivey (2006b), Coupled ROMS-SUNTANS simulations of highly nonlinear internal gravity waves on the Australian northwest shelf, *Proc. 6th Int. Symp. Strat. Flows*, 533-538.

Fringer, O. B., McWilliams, J. C., and R. L. Street (2006c), A new hybrid model for coastal simulations, *Oceanography*, invited, 19 (1), 46-59.

Kanarska, Y., Shchepetkin, A., and J. C. McWilliams (2007), Algorithm for non-hydrostatic dynamics in the Regional Oceanic Modeling System, *Ocean Model.*, 18 (3-4), 143-174.

Leonard, B. P. (2002), Stability of explicit advection schemes. The balance point location rule, *Int. J. Numer. Meth. Fluids*, 38(5), 471-514.

Marchesiello, P., McWilliams, J. C., and A. Shchepetkin (2003), Equilibrium structure and dynamics of the California Current System, *J. Phys. Oceanogr.*, 33 (4), 753-783.

Shchepetkin, A. F., and J. C. McWilliams (2005), The regional oceanic modeling system (ROMS): a split-explicit, free-surface, topography-following-coordinate oceanic model, *Ocean Model.*, 9 (4), 347-404.

Sherwood, C. R., Signell, R. P., Harris, C. K., and B. Butman (2000), Workshop discusses community models for coastal sediment: EOS, *Transactions, American Geophysical Union*, v. 81, no. 43, p. 502.

Staniforth A, and J. Cote (1991), Semi-Lagrangian integration schemes for atmospheric models: a review. *Monthly Weather Review*, 119, 2206–2223.

Wang, B., Fringer, O. B., Giddings, S. N., and D. A. Fong (2008), High-resolution simulations of a macrotidal estuary using SUNTANS, *Ocean Modelling*, in press.

Zhao, G. (2007), Numerical simulation of complex channel flows, Ph.D. Dissertation, Stanford University.

## **HONORS/AWARDS/PRIZES**

Oliver B. Fringer, ONR Young Investigator Award, 2008.